

Accepted Manuscript

Possible effect of hala extract (*Pandanus tectorius*) on immune status, anti-tumour and resistance to *Yersinia ruckeri* infection in rainbow trout (*Oncorhynchus mykiss*)

Elham Awad, Dawn Austin, Alastair Lyndon, Amani Awaad



PII: S1050-4648(19)30086-5

DOI: <https://doi.org/10.1016/j.fsi.2019.02.012>

Reference: YFSIM 5914

To appear in: *Fish and Shellfish Immunology*

Received Date: 10 December 2018

Revised Date: 6 February 2019

Accepted Date: 8 February 2019

Please cite this article as: Awad E, Austin D, Lyndon A, Awaad A, Possible effect of hala extract (*Pandanus tectorius*) on immune status, anti-tumour and resistance to *Yersinia ruckeri* infection in rainbow trout (*Oncorhynchus mykiss*), *Fish and Shellfish Immunology* (2019), doi: <https://doi.org/10.1016/j.fsi.2019.02.012>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Possible effect of hala extract (*Pandanus tectorius*) on immune status, anti-**
2 **tumour and resistance to *Yersinia ruckeri* infection in rainbow trout**
3 **(*Oncorhynchus mykiss*)**

4
5
6
7 Elham Awad^{a,b*}, Dawn Austin^a, Alastair Lyndon^a, Amani Awaad^c.

8
9
10 ^{a,b} Institute of Life and Earth Sciences, Heriot-Watt University, Edinburgh, UK

11 ^b Department of Hydrobiology, National Research Center, Cairo, Egypt

12 ^c Pharmacognosy Department, College of Pharmacy, Salman Bin Abdulaziz University, Al-
13 Kharj, Saudi Arabia

14
15
16 ***Corresponding author:** E. Awad, Department of Hydrobiology, National Research Centre, Cairo,
17 Egypt.

18 E-mail address: elhamsawad@yahoo.com

19 Tel.: +20-1003060046

20 Fax: +20-233370931

21

22 **Abstract:**

23 The possible effect of dietary administration of hala extract (*Pandanus tectorius*) on rainbow
24 trout (*Oncorhynchus mykiss*) immune status as well as its effect as an anti-tumour agent was
25 studied. Fish were divided into 4 groups before feeding with commercial diet (0%, control;
26 0.5%, 1% and 2% of hala extract) for 2 weeks. The effect of diet on the humoral immune
27 parameters, ie total protein, myeloperoxidase content, antiproteases, lysozyme and
28 bactericidal activities were studied. Also, the effect of the diets on the expression of some
29 immune-related genes in rainbow trout head-kidney (*TNF*, *LYZ2*, *IL-8* and *CD-4*) as well as
30 tumour suppressor gene (*WT-1a*) was investigated. At the end of the feeding trial fish groups
31 were challenged with *Yersinia ruckeri*. The results demonstrated enhancement in all the
32 immune parameters in fish fed hala extract diets compared to control fish especially with the
33 highest dose (2%) which recorded the highest significant increase ($p < 0.05$) in some
34 parameters (total protein, myeloperoxidase content, antiproteases, and bactericidal activities)
35 compared to the control. The results obtained from challenge with *Y. ruckeri* revealed
36 reduction in the mortalities in fish groups fed with 1% and 2% doses of hala extract. Feeding
37 with hala extract provoked upregulation in all immune- related genes. Again, the highest dose
38 of hala extract showed a significant upregulation in *WT1a* expression ($p < 0.05$). The current
39 study suggest that the hala extract, especially the highest dose, could be considered a good
40 food additive to improve the immune status, resist tumour formation and to resist or control
41 infectious diseases of rainbow trout.

42

43

44 **Key words:**45 Rainbow trout, *Pandanus tectorius* , immune response, *Yersinia ruckeri*, anti-tumour.

46 1. Introduction:

47 Aquaculture is considered to provide a valuable source of essential protein required for
48 human health. However, the intensive and extensive aquaculture industry is subject to disease
49 outbreaks [1]. However, controlling fish diseases by antibiotics and chemotherapeutics has
50 caused the development of drug resistant pathogens in addition to accumulation of residues in
51 environment and fish tissue and subsequently in humans [2]. On the other hand, medicinal
52 plants provide a promising, alternative method for resisting and/or controlling fish diseases
53 [3]. The world tends to use medicinal plants for treatment not only because they are cost
54 effective, biodegradable, and safe but also for long lasting effects than the synthetic drugs
55 provide and which have faster recovery rates [4]. It is worth mentioning that many medicinal
56 plants have antioxidant properties which delay or prevent oxidative damage, therefore
57 playing a vital role in disease prevention. [5].

58 Previous studies showed a marked enhancement of the fish immune system after
59 administration of different parts of medicinal plants (roots, leaves, seeds, and flowers).
60 Moreover, various levels of immune response depended upon plant concentrations, time and
61 method of administration [3]. For example, dietary supplement with three doses of fenugreek
62 seeds improved the immune status of gilthead seabream (*Sparus aurata* L.) especially with
63 the highest dose (10%) [6]. Moreover, common carp (*Cyprinus carpio*) showed enhancement
64 in immune parameters after administration of a diet supplemented with a 2% dose of *Achillea*
65 *wilhelmsii* leaf extract [7]. Interestingly, catfish (*Clarias gariepinus*) injected with 50 mg/kg
66 of leek leaf extract showed an increase in humoral immune response one month post-
67 injection [8].

68

69 Hala tree (*Pandanus tectorius*) belongs to the family Pandanaceae that comprises around 600
70 members. [9]. This tree was initially cultured in Asia and extends to tropical northern
71 Australia and Pacific islands of Oceania [10]. It contains triterpenoids and flavonoids [11],
72 thus it was successfully used in folk medicine in many countries. In Kiribati, the leaves are
73 used as therapy for cold, influenza, asthma, hepatitis, boils and cancer, while the roots are
74 used to treat haemorrhoids. Fruits, flowers and aerial roots are used to treat digestive and
75 respiratory disorders in Hawaii. While, In Palau, roots and leaves are used to alleviate
76 stomach cramps and vomiting, respectively [10].

77 Previous studies recorded antioxidant, antibacterial, anticoagulant, anti-inflammatory,
78 hepatoprotective, antidiarrheal, anticonvulsant, diuretic and anti-cancer activities for leaf
79 extract [10, 12]. However, there has been no investigation carried out on the effect of hala
80 leaf extract on the immune system of fish. Thus the current study was carried out to
81 investigate the possible effects of dietary supplement of hala leaf extract on rainbow trout
82 immune response either in serum or in cell by examine the expression of some immune-
83 related genes (*TNF*, *LYZ2*, *IL-8* and *CD-4*) in head kidney as well as study its effect as anti-
84 tumour agent by examination of the expression of *WT-1a* gene (tumour suppressor gene).

85

86 2. Materials and methods:

87 2.1. Preparation of plant extract and diets:

88 Hala (*Pandanus tectorius*) leaves were collected from a local market in Saudi Arabia. About
89 one kilogram of dried powder of hala was extracted using 95% alcohol (Merck, Germany) by
90 percolation till exhaustion (4 X 4 l) and filtered off by filter paper. The combined filtrates of
91 the plant were evaporated under reduced pressure and low temperature using rotator

92 evaporator. The obtained residue (250 g) was used in preparing the diets. Four different
93 concentrations were prepared; commercial diet non-supplemented (0%, control), commercial
94 diet supplemented with 0.5 g (0.5%), 1 g (1 %) and 2 g (2%)/ 100 g of hala extract.

95

96 2.2. Fish, experimental design and sampling:

97 Rainbow trout (*Oncorhynchus mykiss*) of average weight 18 ± 1 g were obtained from a
98 commercial fish farm in Scotland, and acclimatized in aerated free flowing freshwater ($14 \pm$
99 2°C). During acclimatization, fish were fed three times daily with a commercial diet
100 (Biomar). Fish were distributed randomly into 4 groups each with 30 fish (10 per replicate)
101 and fed for 14 days with 0.5 g (0.5%), 1g (1%) and 2 g (2%)/100 g of *hala* extract. Controls
102 were fed with commercial diet only to examine the possible mode of action and effect on
103 immune status. Blood was collected from fish anaesthetised using 3- amino benzoic acid
104 ethyl ester; Sigma-Aldrich, Basingstoke, U.K.) by syringe before transfer to Vacuettes
105 without heparin (Greiner, Stonehouse, U.K.) and left to clot for 2 h at 4°C , prior to
106 centrifugation (1600 g, 25 min, 4°C), and stored at -20°C until use. The fish were then
107 sacrificed using an overdose of the above anaesthetic.

108 2.3. Humoral immune parameters:

109 2.3.1. Lysozyme activity:

110 Serum lysozyme activity was measured according to [13]. Briefly, 60 μL of serum was
111 added to 2 mL of a suspension of *Micrococcus lysodeikticus* (0.2 mg ml^{-1} in a 0.05 M sodium
112 phosphate buffer (pH 6.2) and absorbance was measured at 530 nm after 0.5 and 4.5 min on a
113 spectrophotometer. A unit of lysozyme activity was defined as the sample amount causing a
114 decrease in absorbance of 0.001 min^{-1} .

115 2.3.2. Total protein content

116 Total protein was measured by Bradford assay using bovine serum albumin (BSA) as the
117 standard. Briefly, 2 mg ml⁻¹ solution of BSA was prepared and serial dilutions made with
118 phosphate buffer saline (PBS). Around 20 µl of each dilution was added to 1 ml of Bradford
119 reagent (Sigma-Aldrich) before incubated at room temperature for 15 min. The standard
120 curve was prepared by measuring the absorbance of each sample at 595 nm verses the sample
121 concentration. Serum samples were diluted (1: 100) in PBS before 20 µl of each serum
122 dilution was added to 1 ml of Bradford reagent. After incubation for 15 min, the absorbance
123 of the unknown samples was taken and plotted onto the standard curve to obtain the total
124 protein content for each sample [14].

125 2.3.3 Antiproteases activity

126 The serum anti-trypsin activity was measured according to Lange, Guðmundsdottir [15].
127 Thus, 20 µl of standard trypsin solution (Sigma-Aldrich, 5 mg ml⁻¹) was incubated with 20 µl
128 of serum for 10 min at 22°C. Subsequently, 200 µl of 0.1 M PBS (PH 7.2) and 250 µl of 2%
129 azocasein solution (Sigma-Aldrich, 20 mg ml PBS⁻¹) were added. The mixture was incubated
130 for 1 h at 22°C before stopping with the addition of 500 µl of 10 % (v/v) trichloro acetic acid
131 (TCA). Then, the mixture was incubated for 30 min at 2°C before centrifuging at 6000 x g for
132 5 min. About 100 µl of the supernatant was transferred to a 96 microwell flat bottom plate
133 containing 100 µl of 1 N NaOH well⁻¹. The absorbance was read in the ELISA reader at 410
134 nm. Positive control (100%) was prepared by replacing the serum with buffer. For a negative
135 control, buffer replaced both serum and trypsin. The percentage inhibition of trypsin activity
136 was calculated by comparing with a positive control sample.

137

138 2.3.4. Myeloperoxidase content:

139 The myeloperoxidase content of serum was measured according to [16]. Briefly, 50 µl serum
140 was diluted with 135 µl of Ca⁺² and Mg⁺² free HBSS (Sigma-Aldrich) in flat-bottomed 96-
141 well plates. Then, 50 µl of 20 mM 3,3',5,5'-tetramethylbenzidine hydrochloride (TMB,
142 Sigma-Aldrich) and 5 mM H₂O₂ (Sigma-Aldrich) were added (both substrates of peroxidase).
143 The reaction was stopped by adding 50 µl of 4 M sulphuric acid (H₂SO₄) after 2 min. The
144 absorbance was read at 450 nm by ELISA reader. Blank sample was without serum.

145 2.3.5. Bacterial culture and bactericidal activity:

146 *Yersinia ruckeri* was identified and obtained from Heriot Watt University before inoculation
147 into TSA media (Oxoid) for 48 h at 25 °C. The culture was centrifuged for 10 min at 3000 ×g
148 at 4 °C and the pellet resuspended in 0.9% of saline. The bacterial suspensions were counted
149 using a hemocytometer slide at a magnification of 400× on a light microscope.

150 Serum bactericidal activity was done according to Kajita, Sakai [17] using *Yersinia ruckeri*.
151 Briefly, 100 µl of serum was mixed with 100 µl of bacterial suspension, before incubation for
152 1 h at 25 °C. A blank control was also prepared by replacing serum with sterile PBS. The
153 mixture was then diluted with 0.05 M sodium phosphate buffer, PBS (pH 6.2) at a ratio of
154 1:10. Around 50 µl of mixture was plated onto the nutrient agar plates and incubated for 48 h
155 at 25 °C before the number of colonies was counted.

156 2.4. Challenge with *Yersinia ruckeri*:

157 All challenge experiments were done under a UK Home Office Project License (held by A.R.
158 Lyndon) and a UK Home Office Personal License (held by D. A. Austin) and approved by
159 the Heriot-Watt University Ethics Committee.

160 *Yersinia ruckeri* was grown in nutrient broth (Oxoid) for 24 h at 25°C. The culture was
 161 centrifuged at 3000 x g for 10 min at 4°C, before the supernatants were discarded, and the
 162 pellets resuspended in 0.9% (w/v) saline. The challenge test was carried out on fish by
 163 intraperitoneal injection with 0.1 ml volumes containing 10³ cells/fish, as preliminary
 164 experiments had determined this to be the LD80 for the challenge strain of *Y. ruckeri*.
 165 Mortalities were recorded for up to 10 days [18].

166 2.5. Gene expression by real time PCR:

167 Total RNA was extracted from the head kidney and liver using RNA extraction kit (Applied
 168 biosystem, UK). The quantity of RNA was measure using nanodrop. Around 1 ug of RNA was used
 169 in Reverse transcription by using cDNA kit from Fermentas (York, UK).

170 The expression of *TNF*, *LYZ2*, *IL-8*, *CD-4*, *β-actin*, and *WT-1a* genes (Table 1) in head kidney were
 171 analysed by real-time PCR machine (ABI PRISM 7500 instrument, Applied Biosystems) using SYBR
 172 Green PCR Core Reagents (Applied Biosystems). Mixture (comprised of 10 ml of SYBR Green
 173 supermix, 5 ml of primers (0.6 mM each) and 5 ml of cDNA template) were incubated for 2 min at 95
 174 °C, followed by 40 cycles of 15 s at 95 °C, 1 min at 60 °C, and finally 15 s at 95 °C, 1 min at 60 °C
 175 and 15 s at 95 °C. Gene expression was corrected by the reference gene, β-actin in each sample. The
 176 primers used are shown in Table 1. Gene expression of the samples compared to the controls was
 177 calculated according to the following equation:

$$178 \quad (E_{\text{target}})^{\Delta C_{T \text{ target (control - sample)}}$$

$$179 \quad \text{Ratio} = \frac{\quad}{\quad}$$

$$180 \quad (E_{\text{EF1}\alpha})^{\Delta C_{T \text{ target (control - sample)}}$$

181

182 2.6. Statistical analysis

183 Data were expressed as fold increase (mean \pm standard error, SE), obtained by dividing each sample
184 by the mean control value. Values higher than 1 express an increase while values lower than 1 express
185 a decrease in the indicated gene. Data were analyzed by one-way analysis of variance (ANOVA).
186 When differences were found among treatments, Tukey's test was used to compare means by
187 Minitab statistical software (Minitab, Coventry, UK). Differences were considered significant at P
188 <0.05 .

189 3. Results:

190 3.1. Humoral immune parameters:

191 In Lysozyme activity and Myeloperoxidase content (Fig. 1 & 2), the highest significant values ($p <$
192 0.05) were reported in fish groups fed with 0.5% and 2% of hala extract with respect to the values
193 found in the control group (0%). Although all doses of hala extract showed highly serum total protein
194 compared to the control (0%) none of them showed significant differences compared to the values
195 found in control group (Fig.3). The antiprotease activity (Fig.4) was increased in specimens fed with
196 0.5% or 2% doses of hala extract (compared to control), although only in the group fed 2% dose the
197 increases were statistically significant, compared to the values found in control ($p < 0.05$). Regarding
198 bactericidal activity (Fig. 5), all groups fed with hala extract showed high activity compared to the
199 control group (especially with 2% dose) but without significant differences.

200 3.2. Challenge with *Yersinia ruckeri*

201 Fish groups fed with diets supplemented of hala extract for 2 weeks resulted in reduction in
202 mortalities after challenge with *Y. Ruckeri* (Fig.6). The resistance to *Y. ruckeri* infection was increased
203 in fish groups fed for 2 weeks with 2% and 1% doses of hala extract where the survival percent was
204 26.67% and 21.43 %, respectively, compared to control group (33.3 %). However, the difference was
205 not significant.

206 3.3. Gene expression:

207 The results revealed an increase in the expression of immune related genes (*TNF*, *LYZ2*, *IL-8*
208 and *CD-4*) in the head kidney of rainbow trout after administration of diets enriched with
209 0.5% and 1% of hala extract for 2 weeks compared to the expression recorded in the control
210 group (Fig. 7). Interestingly, the lowest dose (0.5%) showed the highest expression compared
211 to other groups and control group while only being statistically significant ($p < 0.05$) for *TNF*
212 and *IL-8* compared to the control.

213 Moreover, the results showed significant up- regulation in *WT-1a* gene of head kidney ($p <$
214 0.05), in fish fed with 1% and 2% doses of hala extract compared to control. The highest
215 expression was recorded in the dose of 2%.

216 Discussion:

217 Lysozyme is an important non-specific immune parameter which plays a vital role in fish defence
218 mechanisms against diseases. It is responsible for opsonin, and thus activates the complement system
219 and phagocytes [19]. Our results revealed a significant enhancement in serum lysozyme activity of
220 the fish group fed 0.5% and 2% of hala extract, respectively, for 2 weeks compared to the control
221 group (0%). This could be attributed to a dose dependent effect of hala extract on rainbow trout.
222 Similar observation was reported in lysozyme activity of rainbow trout fed for 2 weeks with 0.5% and
223 1% of tetra (*Cotinus coggyria*) [20], and 1% & 2% of black cumin oil (*Nigella sativa*) [21], . Also, the
224 highest activity depended mainly on the dose administration.

225 Myeloperoxidase is an important enzyme expressed mainly in neutrophils, which have the ability to
226 produce hypochlorous acid from one of the oxidative radicals (H_2O_2) [22]. This process has a great
227 benefit to kill invading microorganisms [23]. The results showed an increase in myeloperoxidase
228 content in all treatment groups with hala extract, especially in those fed with 0.5 % and 2% where a
229 significant value ($p < 0.5$) was recorded compared to control group. Similar to our study,
230 myeloperoxidase was increased significantly in rainbow trout fed for two weeks with 1% and 2% of
231 lupin (*Lupinus perennis*), mango (*Mangifera indica*), and nettle (*Urtica dioica*) [3]. Also, the highest

232 myeloperoxidase content and lysozyme activity values were recorded in rainbow trout fed 0.1% and
233 0.5% of caper leaf extract for 4 weeks [19]. It is worth mentioning that time and dosage are two
234 important factors which control the efficiency of plant immunostimulants. For example,
235 Christyapita, Divyagnaneswari [24] noticed a significant increase in common tilapia fed with diets
236 containing different concentrations of false daisy leaf for 1 week, while feeding for 2 or 3 weeks
237 didn't showed any significant increase.

238

239 Serum protein is an important parameter in humeral immune system in fish. Its composition plays a
240 vital role in keeping fish healthy. Moreover, the most important role played by acute phase proteins is
241 in limiting the spread of infectious agents through repairing tissue damage and killing micro-
242 organisms [25, 26]. Present results also demonstrated enhancement in total protein value in all groups
243 that received different concentration of hala extract groups as compared to the control (especially with
244 the dose of 2%). This is in agreement with Dügenci, Arda [27] who reported increases in serum
245 protein levels in rainbow trout fed with 0.1% and 1% of ginger (*Zingiber officinale*), nettle (*Urtica*
246 *dioica*) and mistletoe (*Viscum album*). Several studies reported an increase of serum protein levels in
247 fish species after using dietary supplement with plants as immunostimulants [6, 7, 28, 29]. Moreover,
248 they suggested that elevation in fish total protein were probably a result of enhancement of the non-
249 specific immune response.

250 Antiproteases or protease inhibitors are active molecules in the non-specific immune system that
251 inhibit the action of proteases either by binding to their active sites or by 'trapping' the protease to
252 prevent protein hydrolysis [30] and thus limiting the growth of invading bacteria in fish [31]. In the
253 present study, the highest dose of hala extract recorded the highest significant enhancement in serum
254 antiproteases compared to the control. In agreement with our result several studies recorded
255 enhancement in fish species after administration of diets supplemented with medicinal plants [6, 32,
256 33]. It is worth emphasizing that the increase in fish immune response depend mainly on dose and
257 time of administration. Moreover, the response also depends on fish species. For example, 0.5% dose

258 of garlic reported the highest antiprotease activity in rainbow trout after feeding for 2 weeks [33].
259 Although, the highest antiprotease in Asian seabass fed for 2 weeks was reported at 1.5% dose of
260 garlic [34].

261 Various humoral molecules involved in non-specific immune response have a power to protect the
262 fish from invading microbes [35]. Serum bactericidal activity is a lysin mechanism known for the
263 killing and clearing of pathogenic organisms in fish [31]. In our study, *Y. ruckeri* was used as a model
264 to examine the activity of hala extract to kill the bacterial infection. The strength of immune
265 molecules in fish serum to kill *Y. ruckeri* can be detected by the lowest number of bacterial colonies
266 grown on media. Fish groups fed with hala extract revealed higher bactericidal activity compared to
267 control, especially in the group fed with the highest dose (2%). Similarly, using the highest dose of
268 black cumin seed (3%) as food supplement in rainbow trout diet caused higher bactericidal activity against
269 *Aeromonas hydrophila*, [21]. Also, rohu (*Labeo rohita*) recorded an enhancement in serum
270 bactericidal activity against *A. hydrophila* after feeding for 2 weeks with doses of prickly chaff-flower
271 seed and the activity was elevated with higher concentration of seeds [35].

272 Challenge with target pathogen is one of the most valuable tests to evaluate the efficiency of
273 immunostimulant to resist microbes. The resistance level of fish can be recognized from the survival
274 percent after a bacterial infection [36]. The results revealed that dietary supplement with hala extract
275 relatively increased the resistance of rainbow trout against *Y. ruckeri*, where the highest doses (2%
276 and 1% respectively) recorded the highest resistance. In agreement with our study using stinging
277 nettle (*Urtica dioica*) as food supplement of rainbow trout diet increased the resistance to *Y. ruckeri*,
278 especially with the highest dose [37]. However, some immunostimulant can enhance the resistance of
279 fish against some bacteria but failed to resist the other. For example an improvement in survival
280 percent of rainbow trout fed with probiotic for 2 weeks was recorded following challenge with
281 *Aeromonas salmonicida*, *Vibrio ordalii*, and *Y. ruckeri*, but not so with *V. anguillarum* [38].
282 Previous studies showed reduction in mortality against bacterial infections in fish after using plant
283 immunostimulant [3, 39, 40].

284

285 *TNF- α* , is one of pro-inflammatory cytokines, that mainly produced by activated
286 monocytes/macrophages and regulate the expression of many cytokines [41]. Many studies revealed
287 that using plant immunostimulants in fish can induce pro-inflammatory responses. For example, using
288 0.1% of caper as supplement in rainbow trout diet caused up-regulation in the expression of *TNF- α* of
289 head kidney [42]. Also, an up-regulation in *TNF- α* expression was observed in common carp treated
290 with *Rehmannia glutinosa* in spleen, head kidney and gut [43]. Moreover, common carp fed with
291 different concentrations of guava leaf powder showed an increase in *TNF- α* expression in the head-
292 kidney, hepatopancreas, and intestine [44]. Similarly our study demonstrated up-regulation of *TNF- α*
293 expression of head kidney in fish groups fed with hala extract compared to control. Increasing in the
294 *TNF- α* levels could be attributed to the activity of the compounds in hala extract like flavonoids and
295 antioxidant [12].

296 *IL-8* is another pro-inflammatory cytokine, that produced in response to many stimulation factors like
297 cytokines, LPS and viruses [45]. It plays an important role in attract T-lymphocytes and neutrophils to
298 sites of inflammation [46]. Similar to *TNF- α* expression, result demonstrated an increase in *IL-8* in
299 rainbow trout head kidney in groups received 0.5% and 1 % doses of hala extract. Although, only 0.5%
300 dose recorded significant difference with the control. In agreement with this study, *IL-8* expression
301 increased in rainbow trout head kidney after fed diets supplemented with 1% and 2% of stinging
302 nettle [47] and 0.1% of caper [42]. Interestingly, *IL-8* expression in rainbow trout spleen was
303 unaffected by the green tea supplementation, while in head kidney the expression showed significant
304 increase especially in fish fed with 500 mg kg⁻¹ of green tea [48]. Similarly, gilthead seabream fed
305 dietary supplement with 10% fenugreek showed enhancement in *IL-8* expression of head kidney after
306 4 weeks [6].

307 *CD4* is an important co-receptor has been reported in teleosts, expressed on T- helper cell,
308 monocytes and macrophages [49]. *CD4* is binding to major histocompatibility complex (MHC) class I
309 molecules on the antigen-presenting cells, stabilizing the interaction between the T cell receptor

310 complex (TCR) and the MHC [50]. The results demonstrated an increase in *CD4* expression in head
311 kidney in group fed with hala extract compared to control group. Similar observations have been in
312 other fish species after administration immunostimulant. For example; Atlantic salmon (*Salmo salar*)
313 injected with lipopolysaccharide (LPS) and β -glucan as immunostimulant, showed an enhancement in
314 the expression of *CD4* in head kidney [51]. Also, an increase in *CD4* expression of posterior intestine
315 of European sea bass (*Dicentrarchus labrax*) has been reported after feeding diets contain low level of
316 synbiotic additive (mannan oligosaccharides) [52]. The slightly higher expression values of *CD4* of
317 fish group fed with hala extract may indicate an early adaptive immune response.

318 Lysozyme gene is a bactericidal enzyme that mainly present in lymphoid tissue like head kidney and
319 thymus as well as serum, mucus, gills [53]. There is two types of lysozyme (types I and II), have been
320 identified in the kidney of rainbow trout. Particular, lysozyme type II (*LYZ2*) showed a potential
321 antibacterial activity against four gram-negative bacteria. Such finding supports the role which
322 lysozyme plays in non-specific immune defence in fish [54]. Our study reported the highest level of
323 *LYZ2* expression in head kidney of fish group fed with the lowest dose of hala extract (0.5%), this is
324 agreement with the result obtained from analysis serum lysozyme level for the same group. Similar
325 observation have been reported in common carp fed for 8 weeks with diet supplements with date palm
326 fruit (200 ml kg⁻¹) where showed a remarkable increase in the expression level of lysozyme gene in
327 head kidney (*LYZ2*) and serum lysozyme compared to control [55].

328 The kidney is plays important functions in fish, not only in immune system for production the
329 leucocytes but also as osmoregulatory function [56]. Thus any malfunction in this organ in really
330 preferable. Wilms' tumour suppressor *WT1* is a tumor suppressor gene, any mutation can led to
331 Wilms' tumour, a pediatric kidney cancer [57]. *WT1* is a modulatory gene involved in cell growth and
332 development of the urogenital system development (kidney and gonad) [58]. *WT1* have been
333 identified in fish species [59-61]. The suppression of *WT1* led to appear the edema in zebrafish which
334 suggest that it is involved in pronephros development [62]. *WT1a* and *WT1b* are two types have been
335 reported from *WT1* in zebrafish. Inactivation of *wt1a* leads to the absence of glomeruli while targeting
336 of *wt1b* resulted in the formation of renal cysts [60]. The result showed an increase in *WT1a*

337 expression in head kidney of fish group treatment with hala extract, especially in the group fed with
338 2% which recorded the highest value. This could be contributed to the activity of ROS in plant extract
339 compounds to activate the tumour suppressor gene and suggest the role of this plant to work as anti-
340 tumour agent in kidney.

341 In conclusion, our results demonstrated that dietary supplement with hala extract to rainbow trout for
342 two weeks stimulates the non-specific immune response and increase its resistance toward *Y. ruckeri*
343 infection. The current results suggest that the hala extract, specially the highest dosage (2%) could be
344 considered as a good fish food supplements to enhance the immune system and resist and/or control
345 pathogenic bacterial, in addition to its potential effect to resist the tumour formulation. Future
346 investigations could focus on the effects of long feeding time on the immune status and general health
347 status of fish.

348 References:

- 349 1. Bondad-Reantaso MG, Subasinghe RP, Arthur JR, Ogawa K, Chinabut S, Adlard R, et al.
350 Disease and health management in Asian aquaculture. *Veterinary parasitology*. 2005 132:249-72.
- 351 2. Alexander CP, John Wesly Kirubakaran C, Michael RD. Water soluble fraction of *Tinospora*
352 *cordifolia* leaves enhanced the non-specific immune mechanisms and disease resistance in
353 *Oreochromis mossambicus*. *Fish Shellfish Immunol*. 2010 29:765-72.
- 354 3. Awad E, Awaad A. Role of medicinal plants on growth performance and immune status in
355 fish. *Fish Shellfish Immunol*. 2017 67:40-54.
- 356 4. Wu C-C, Liu C-H, Chang Y-P, Hsieh S-L. Effects of hot-water extract of *Toona sinensis* on
357 immune response and resistance to *Aeromonas hydrophila* in *Oreochromis mossambicus*. *Fish*
358 *Shellfish Immunol*. 2010 29:258-63.
- 359 5. Hudec J, Burdová M, Kobida Lu, Komora L, Macho V, Kogan G, et al. Antioxidant capacity
360 changes and phenolic profile of *Echinacea purpurea*, nettle (*Urtica dioica* L.), and dandelion
361 (*Taraxacum officinale*) after application of polyamine and phenolic biosynthesis regulators. *Journal*
362 *of agricultural and food chemistry*. 2007 55:5689-96.
- 363 6. Awad E, Cerezuela R, Esteban MÁ. Effects of fenugreek (*Trigonella foenum graecum*) on
364 gilthead seabream (*Sparus aurata* L.) immune status and growth performance. *Fish Shellfish*
365 *Immunol*. 2015 45:454-64.
- 366 7. Adel Milad, Mehdi Gholaghaie, Mohammad Binaii, Poria Khanjany, Awad E. Effect of dietary
367 *Achillea wilhelmsii* extract on growth performance, and immune status of common carp (*Cyprinus*
368 *carpio*). *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2016 7:1037- 46.
- 369 8. Abbas WT, Awad E. Effect of leek (*Allium ampeloprasum* L.) extract on biotransformation
370 enzymes and innate immunity of catfish (*Clarias gariepinus*) exposed to Benzo [a] Pyrene. *Research*
371 *Journal of Pharmaceutical, Biological and Chemical Sciences*. 2016 7:2211-24.
- 372 9. Englberger L, Aalbersberg W, Fitzgerald MH, Marks GC, Chand K. Provitamin A carotenoid
373 content of different cultivars of edible pandanus fruit. *Journal of food composition and analysis*.
374 2003 16:237-47.

- 375 10. Baba S, Chan HT, Kezuka M, Inoue T, Chan EWC. *Artocarpus altilis* and *Pandanus tectorius*:
376 Two important fruits of Oceania with medicinal values. Emirates Journal of Food and Agriculture.
377 2016:531-9.
- 378 11. Gupta V, Niazi J, Kehal JK, Bansal P. Anti-Inflammatory and Anti-Nociceptive Activity of
379 *Pandanus tectorius* Parkinson. Research Journal of Pharmacognosy and Phytochemistry. 2010 2:193-
380 5.
- 381 12. Omodamiro O, Ikekamma C. In vitro study of antioxidant and anticoagulant activities of
382 ethanol extract of *Pandanus tectorius* leaves. International Blood Research Reviews. 2016 5:1-11.
- 383 13. Parry RM, Chandan RC, Shahani KM. A rapid and sensitive assay of muramidase.
384 Proceedings of the Society for Experimental Biology and Medicine Society for Experimental Biology
385 and Medicine (New York, NY): Royal Society of Medicine; 1965, p. 384-6.
- 386 14. Bradford MM. Foreign Patent Documents. Biochemistry. 1976 72:248-54.
- 387 15. Lange S, Guðmundsdóttir BK, Magnadóttir B. Humoral immune parameters of cultured
388 Atlantic halibut (*Hippoglossus hippoglossus* L.). Fish Shellfish Immunol. 2001 11:523-35.
- 389 16. Quade MJ, Roth JA. A rapid, direct assay to measure degranulation of bovine neutrophil
390 primary granules. Vet Immunol Immunopathol. 1997 58:239-48.
- 391 17. Kajita Y, Sakai M, Atsuta S, Kobayashi M. The immunomodulatory effects of levamisole on
392 rainbow trout, *Oncorhynchus mykiss*. Fish Pathology. 1990 25:93-8.
- 393 18. Austin B, Austin DA. Bacterial fish pathogens : disease of farmed and wild fish. 4th ed.
394 Dordrecht Chichester: Springer ; Published in association with Praxis Pub.; 2007.
- 395 19. Kumari J, Sahoo PK. Dietary levamisole modulates the immune response and disease
396 resistance of Asian catfish *Clarias batrachus* (Linnaeus). Aquaculture Research. 2006 37:500-9.
- 397 20. Bilen S, Bulut M, Bilen AM. Immunostimulant effects of *Cotinus coggyria* on rainbow trout (
398 *Oncorhynchus mykiss*). Fish Shellfish Immunol. 2011 30:451-5.
- 399 21. Awad E, Austin D, Lyndon AR. Effect of black cumin seed oil (*Nigella sativa*) and nettle
400 extract (Quercetin) on enhancement of immunity in rainbow trout, *Oncorhynchus mykiss*(Walbaum).
401 Aquaculture. 2013 388:193-7.
- 402 22. Dalmo R, Ingebrigtsen K, Børgwald J. Non-specific defence mechanisms in fish, with particular
403 reference to the reticuloendothelial system (RES). J Fish Dis. 1997 20:241-73.
- 404 23. Johnston Jr R. Oxygen metabolism and the microbicidal activity of macrophages. Federation
405 proceedings; 1978, p. 2759-64.
- 406 24. Christyapita D, Divyagnaneswari M, Dinakaran Michael R. Oral administration of *Eclipta*
407 *alba* leaf aqueous extract enhances the non-specific immune responses and disease resistance of
408 *Oreochromis mossambicus*. Fish and shellfish immunology. 2007 23:840-52.
- 409 25. Larsen MH, Larsen JL, Olsen JE. Chemotaxis of *Vibrio anguillarum* to fish mucus: role of the
410 origin of the fish mucus, the fish species and the serogroup of the pathogen. FEMS Microbiology
411 Ecology. 2001 38:77-80.
- 412 26. Gerwick L, Steinhauer R, Lapatra S, Sandell T, Ortuno J, Hajiseyedjavadi N, et al. The acute
413 phase response of rainbow trout (*Oncorhynchus mykiss*) plasma proteins to viral, bacterial and
414 fungal inflammatory agents. Fish Shellfish Immunol. 2002 12:229-42.
- 415 27. Dügenci SK, Arda N, Candan A. Some medicinal plants as immunostimulant for fish. Journal
416 of Ethnopharmacology. 2003 88:99-106.
- 417 28. Awad E, Austin B. Use of lupin, *Lupinus perennis*, mango, *Mangifera indica*, and stinging
418 nettle, *Urtica dioica*, as feed additives to prevent *Aeromonas hydrophila* infection in rainbow trout,
419 *Oncorhynchus mykiss* (Walbaum). J Fish Dis. 2010 33:413-20.
- 420 29. Dorucu M, Colak SO, Ispir U, Altinterim B, Celayir Y. The effect of black cumin seeds (*Nigella*
421 *sativa*) on the immune response of rainbow trout (*Oncorhynchus mykiss*). Proceedings of the Middle
422 East & North Africa (MENA) Region Animal Wealth Research Conference Future of animal wealth,
423 Cairo International Convention Center, Egypt, 16-18 October, 2008: Massive Conferences and Trade
424 Fairs; 2008, p. 543-50.

- 425 30. Laskowski Jr M, Kato I. Protein inhibitors of proteinases. Annual review of biochemistry.
426 1980 49:593-626.
- 427 31. Ellis A. Innate host defense mechanisms of fish against viruses and bacteria. Developmental
428 & Comparative Immunology. 2001 25:827-39.
- 429 32. Awad ES. Studies on plant based dietary supplements for control of *Aeromonas hydrophila*
430 infections in rainbow trout (*Oncorhynchus mykiss* Walbaum): Heriot-Watt University; 2010.
- 431 33. Nya EJ, Austin B. Use of garlic, *Allium sativum*, to control *Aeromonas hydrophila* infection in
432 rainbow trout, *Oncorhynchus mykiss* (Walbaum). J Fish Dis. 2009 32:963-70.
- 433 34. Talpur AD, Ikhwanuddin M. Dietary effects of garlic (*Allium sativum*) on haemato-
434 immunological parameters, survival, growth, and disease resistance against *Vibrio harveyi* infection
435 in Asian sea bass, *Lates calcarifer* (Bloch). Aquaculture. 2012 364:6-12.
- 436 35. Rao Y, Das B, Jyotirmayee P, Chakrabarti R. Effect of *Achyranthes aspera* on the immunity
437 and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. Fish and shellfish immunology.
438 2006 20:263-73.
- 439 36. Wassom D, Kelly E. The role of the major histocompatibility complex in resistance to parasite
440 infections. Critical Reviews in Immunology. 1990 10:31-52.
- 441 37. Adel M, Caipang CMA, Dawood MA. Immunological responses and disease resistance of
442 rainbow trout (*Oncorhynchus mykiss*) juveniles following dietary administration of stinging nettle
443 (*Urtica dioica*). Fish Shellfish Immunol. 2017 71:230-8.
- 444 38. Robertson P, O'Dowd C, Burrells C, Williams P, Austin B. Use of *Carnobacterium* sp. as a
445 probiotic for Atlantic salmon (*Salmo salar* L.) and rainbow trout (*Oncorhynchus mykiss*, Walbaum).
446 Aquaculture. 2000 185:235-43.
- 447 39. Baba E, Acar Ü, Yilmaz S, Zemheri F, Ergün S. Dietary olive leaf (*Olea europea* L.) extract
448 alters some immune gene expression levels and disease resistance to *Yersinia ruckeri* infection in
449 rainbow trout *Oncorhynchus mykiss*. Fish Shellfish Immunol. 2018 79:28-33.
- 450 40. Nya EJ, Austin B. Development of immunity in rainbow trout (*Oncorhynchus mykiss*,
451 Walbaum) to *Aeromonas hydrophila* after the dietary application of garlic. Fish Shellfish Immunol.
452 2011 30:845-50.
- 453 41. Zou J, Peddie S, Scapigliati G, Zhang Y, Bols N, Ellis A, et al. Functional characterisation of the
454 recombinant tumor necrosis factors in rainbow trout, *Oncorhynchus mykiss*. Developmental &
455 Comparative Immunology. 2003 27:813-22.
- 456 42. Bilen S, Altunoglu YC, Ulu F, Biswas G. Innate immune and growth promoting responses to
457 caper (*Capparis spinosa*) extract in rainbow trout (*Oncorhynchus mykiss*). Fish Shellfish Immunol.
458 2016 57:206-12.
- 459 43. Wang J-L, Meng X-l, Lu R-h, Wu C, Luo Y-T, Yan X, et al. Effects of *Rehmannia glutinosa* on
460 growth performance, immunological parameters and disease resistance to *Aeromonas hydrophila* in
461 common carp (*Cyprinus carpio* L.). Aquaculture. 2015 435:293-300.
- 462 44. Giri SS, Sen SS, Chi C, Kim HJ, Yun S, Park SC, et al. Effect of guava leaves on the growth
463 performance and cytokine gene expression of *Labeo rohita* and its susceptibility to *Aeromonas*
464 *hydrophila* infection. Fish Shellfish Immunol. 2015 46:217-24.
- 465 45. Seppola M, Larsen AN, Steiro K, Robertsen B, Jensen I. Characterisation and expression
466 analysis of the interleukin genes, IL-1 β , IL-8 and IL-10, in Atlantic cod (*Gadus morhua* L.). Molecular
467 immunology. 2008 45:887-97.
- 468 46. Mukaida N, Harada A, Matsushima K. Interleukin-8 (IL-8) and monocyte chemotactic and
469 activating factor (MCAF/MCP-1), chemokines essentially involved in inflammatory and immune
470 reactions. Cytokine & growth factor reviews. 1998 9:9-23.
- 471 47. Awad E, Mitchell WJ, Austin B. Effect of dietary supplements on cytokine gene expression in
472 rainbow trout, *Oncorhynchus mykiss* (Walbaum). J Fish Dis. 2011 34:629-34.
- 473 48. Nootash S, Sheikhzadeh N, Baradaran B, Oushani AK, Moghadam MRM, Nofouzi K, et al.
474 Green tea (*Camellia sinensis*) administration induces expression of immune relevant genes and

- 475 biochemical parameters in rainbow trout (*Oncorhynchus mykiss*). Fish Shellfish Immunol. 2013
476 35:1916-23.
- 477 49. Suetake H, Araki K, Suzuki Y. Cloning, expression, and characterization of fugu CD4, the first
478 ectothermic animal CD4. Immunogenetics. 2004 56:368-74.
- 479 50. Moore L, Somamoto T, Lie K, Dijkstra J, Hordvik I. Characterisation of salmon and trout CD8 α
480 and CD8 β . Molecular immunology. 2005 42:1225-34.
- 481 51. Hølvold LB. Immunostimulants connecting innate and adaptive immunity in Atlantic salmon
482 (*Salmo salar*): Universitetet i Tromsø; 2007.
- 483 52. Torrecillas S, Rivero-Ramírez F, Izquierdo M, Caballero M, Makol A, Suarez-Bregua P, et al.
484 Feeding European sea bass (*Dicentrarchus labrax*) juveniles with a functional synbiotic additive
485 (mannan oligosaccharides and *Pediococcus acidilactici*): An effective tool to reduce low fishmeal and
486 fish oil gut health effects? Fish Shellfish Immunol. 2018 81:10-20.
- 487 53. Magnadóttir B, Lange S, Gudmundsdóttir S, Bøggwald J, Dalmo R. Ontogeny of humoral
488 immune parameters in fish. Fish Shellfish Immunol. 2005 19:429-39.
- 489 54. Grinde B. Lysozyme from rainbow trout, *Salmo gairdneri* Richardson, as an antibacterial
490 agent against fish pathogens. J Fish Dis. 1989 12:95-104.
- 491 55. Hoseinifar SH, Khalili M, Rufchaei R, Raeisi M, Attar M, Cordero H, et al. Effects of date palm
492 fruit extracts on skin mucosal immunity, immune related genes expression and growth performance
493 of common carp (*Cyprinus carpio*) fry. Fish Shellfish Immunol. 2015 47:706-11.
- 494 56. Henderson I, Hazon N, Hughes K. Hormones, ionic regulation and kidney function in fishes.
495 Symposia of the Society for Experimental Biology; 1985, p. 245-65.
- 496 57. Call KM, Glaser T, Ito CY, Buckler AJ, Pelletier J, Haber DA, et al. Isolation and
497 characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus.
498 cell. 1990 60:509-20.
- 499 58. Kreidberg JA, Sariola H, Loring JM, Maeda M, Pelletier J, Housman D, et al. WT-1 is required
500 for early kidney development. cell. 1993 74:679-91.
- 501 59. Klüver N, Herpin A, Braasch I, Drießle J, Scharfl M. Regulatory back-up circuit of medaka Wt1
502 co-orthologs ensures PGC maintenance. Developmental biology. 2009 325:179-88.
- 503 60. Perner B, Englert C, Bollig F. The Wilms tumor genes wt1a and wt1b control different steps
504 during formation of the zebrafish pronephros. Developmental biology. 2007 309:87-96.
- 505 61. Jiang D, Chen J, Fan Z, Tan D, Zhao J, Shi H, et al. CRISPR/Cas9-induced disruption of wt1a
506 and wt1b reveals their different roles in kidney and gonad development in Nile tilapia.
507 Developmental biology. 2017 428:63-73.
- 508 62. Hsu H-J, Lin G, Chung B-c. Parallel early development of zebrafish interrenal glands and
509 pronephros: differential control by wt1 and ff1b. Development. 2003 130:2107-16.

510

511

512

513

514

515

ACCEPTED MANUSCRIPT

Table 1. Primers used for real-time PCR.

Gene name	Primer sequences (5' - 3')	Products size	GenBank number
Cluster of differentiation (CD4)	GCCCTGCAGAGGACAAATCT TACAAAGGCCACTGGAGCTG	171	NM_001124539.1
Lysozyme II (LYZ2)	TCCAGATCAACAGCCGCTAC GATTCCGTTCCGGTCCAACA	149	NM_001124716.1
Interleukin 8 receptor (IL-8)	CGGTGCCGTCATATTCCTGT GGGTCAGGGACTGTTGACTG	110	NM_001124279.1
Beta-actin (β -actin)	ATGGGCCAGAAAGACAGCTACGTG CTTCTCCATGTCGTCCCAGTTGGT	186	AJ438158.1
Tumor necrosis factor TNF	CAAGAGTTTGAACCTCATTAG GCTGCTGCCGCACATAAAG	130	NM_001124374
Wilms' tumor suppressor 1a (WT-1a)	ATGTTTCCAGCAACGCACCCTA GAACTGGGAGGAGTGGTGTG	129	NM_001124294.1




Figure 1. Serum lysozyme activity of rainbow trout fed for 2 weeks with dietary hala extract supplemented doses: 0% (control), 0.5%, 1% and 2%. Data are presented as mean \pm S.E. Asterisk represents significant difference from control $p \leq 0.05$. Bars = mean \pm S.E.




Figure 2. Serum Myeloperoxidase content of rainbow trout fed for 2 weeks with dietary hala extract supplemented doses: 0% (control), 0.5%, 1% and 2%. Data are presented as mean \pm S.E. Asterisk represents significant difference from control $p \leq 0.05$. Bars = mean \pm S.E.




Figure 3. Serum total protein of rainbow trout fed for 2 weeks with dietary hala extract supplemented doses: 0% (control), 0.5%, 1% and 2%. Data are presented as mean \pm S.E. Asterisk represents significant difference from control $p \leq 0.05$. Bars =mean \pm S.E.




Figure 4. Serum Antiprotease of rainbow trout fed for 2 weeks with dietary hala extract supplemented doses: 0% (control), 0.5%, 1% and 2%. Data are presented as mean \pm S.E. Asterisk represents significant difference from control $p \leq 0.05$. Bars =mean \pm S.E.




Figure 5. Serum bactericidal activity of rainbow trout fed for 2 weeks with dietary hala extract supplemented doses: 0% (control), 0.5%, 1% and 2%. Data are presented as mean \pm S.E. Asterisk represents significant difference from control $p \leq 0.05$. Bars =mean \pm S.E.




Figure 6. Survival of rainbow trout fed for 2 weeks with dietary hala extract supplemented doses: 0% (control), 0.5%, 1% and 2% followed by challenge with *Yersinia ruckeri*. Data are presented as mean \pm S.E. Asterisk represents significant difference from control $p \leq 0.05$. Bars = mean \pm S.E.




Figure 7. Expression of immune related genes (TNF, LYZ2, IL-8 and CD-4) and tumour suppressor genes (WT-1a) in the head kidney of rainbow trout fed dietary hala extract supplemented doses of 0.5%, 1% and 2% for 2 weeks. Data are expressed as fold increase (mean \pm standard error, SE), obtained by dividing each sample value by the mean control value at the same sampling time. Values higher than 1 express an increase while values lower than 1 express a decrease in the indicated gene. Asterisks denote significant differences between control and treatment groups ($P < 0.05$).

- Effects of hala extract (*Pandanus tectorius*) on the immune status of rainbow trout
- Hala extract provoked significant up-regulation in most of immune-related.
- Hala extract can use to resist tumour formation.
- Feeding with hala extract increased resistance toward *Yersinia ruckeri* infection

ACCEPTED MANUSCRIPT